

Flu Decision Chart

Protecting Yourself, Protecting Others

IF YOU HAVE FLU SYMPTOMS

Use the guidelines in this table to help make the best decision for you and your loved ones. Always use hygiene and prevention measures to avoid contamination:

- > Wash your hands frequently.
- > Cough or sneeze into a tissue you can dispose of or your sleeve
- > Keep your surroundings clean.

<p>SITUATION FOR AN ADULT OR CHILD The person does not have a fever (temperature less than 38°C or 100.4°F), but does have these symptoms: > Sore throat > Stuffy nose > Runny nose > Cough</p>	<p>DECISION Probably a cold. Rest is indicated.</p>
<p>SITUATION FOR AN ADULT OR CHILD The person has a fever over 38°C (100.4°F). The fever came on suddenly and is accompanied by these symptoms: > Cough > Sore throat > Significant fatigue > Headache > Muscle aches</p>	<p>DECISION Probably the flu. Rest at home until 24 hours after the fever is gone. Drink plenty of liquids and take fever reducers such as ibuprofen and acetaminophen. Children with fever should not take aspirin.</p>
<p>SITUATION FOR AN ADULT OR CHILD AT RISK OF COMPLICATIONS The person has a fever over 38°C (100.4°F) and belongs to a group at risk of developing complications (children under 2 years of age, the elderly, pregnant women, and individuals with chronic diseases).</p>	<p>DECISION See a doctor today.</p>
<p>SITUATION FOR AN ADULT OR CHILD The person has a fever and one of these symptoms: > Shortness of breath > Difficulty breathing > Painful breathing > Vomiting for more than four hours > Fever in a child who is too quiet and less active than normally or who refuses to play or is agitated</p>	<p>DECISION Go to the emergency room immediately. Call 9-1-1, if necessary.</p>
<p>SITUATION FOR AN ADULT OR CHILD The person has a fever over 38°C (100.4°F) and one of the following: > Difficulty breathing that persists or worsens > Blue lips > Difficulty moving > Severe neck stiffness > Drowsiness, confusion, disorientation, or difficulty being roused > Convulsions > No urination for 12 hours > Fever in an infant under 3 months old</p>	<p>DECISION Go to the emergency room immediately. Call 9-1-1, if necessary.</p>



This publication was supported by Grant # 1H75TP000364-01 from the CDC.